

Transfer of two species of *Aphanes* L. to the genus *Alchemilla* L. (Rosaceae)

Agustín Lahora

Department of Biology and Geology, University of Almería, 04120-Almería, Spain.

Author for correspondence: agustinla@gmail.com

Received: 12 octubre 2024. **Accepted:** 23 octubre 2024. **Published on line:** 28 octubre 2024.

Edited by: Carlos M. Romero Zarco

Abstract

Recent molecular phylogeny studies have shown that the least disruptive taxonomic option in the tribe *Potentilleae* Sweet is to include the taxa of the genera *Aphanes* L. and *Lachemilla* (Focke) Rydb. in the genus *Alchemilla* L. Two combinations are made and incorrect attributions are rectified.

Key words: Rosaceae, *Potentilleae*, *Alchemilla*, *Aphanes*.

Resumen

Transferencia de dos especies de *Aphanes* L. al género *Alchemilla* L. (Rosaceae).

Recientes estudios de filogenia molecular han puesto de manifiesto que la opción taxonómica menos disruptiva en la tribu *Potentilleae* Sweet es incluir los taxones de los géneros *Aphanes* L. y *Lachemilla* (Focke) Rydb. en el género *Alchemilla* L. Se realizan dos combinaciones y se rectifican unas atribuciones incorrectas.

Palabras clave: Rosaceae, *Potentilleae*, *Alchemilla*, *Aphanes*.

Molecular phylogeny studies of the family Rosaceae have shown that *Alchemilla* L. is a polyphyletic genus and extending its circumscription to include *Aphanes* L. and *Lachemilla* (Focke) Rydb. is desirable (Potter *et al.*, 2007; Gehrke *et al.*, 2008). This option, which had previously been adopted by some authors (Murbeck, 1915; Jahandiez & Maire, 1932; Rothmaler, 1935; Maire, 1980; Stevens, 2001; Kalkman, 2004), has now been widely accepted (POWO, 2024; WFO, 2024). Other proposals, such as the creation of the supergenus *Potentilla* L. (Christenhusz & Väre, 2012), have been considered more disruptive (Dobeš & Paule, 2010; Eriksson *et al.*, 2022).

This paper proposes combinations for the transfer of two species of *Aphanes* to the genus *Alchemilla*. Incorrect attributions are also rectified.

***Alchemilla lusitanica* (Frost-Olsen) Lahora, comb. nov.**

Basionym: \equiv *Aphanes lusitanica* Frost-Olsen in Anales Jard. Bot. Madrid 55(1): 195 (1997).

Aphanes lusitanica was described by Frost-Olsen (1997) during the revision of the genus for *Flora iberica* (Frost-Olsen, 1998). It is a Portuguese endemism with a very restricted distribution that had been overlooked in previous studies of the genus (Walters, 1968; Lippert, 1984).

***Alchemilla maroccana* (Hyl. & Rothm.) Lahora, comb. nov.**

Basionym: \equiv *Aphanes maroccana* Hyl. & Rothm. in Svensk Bot. Tidskr. 32(2): 188, f. 2 (1938).

Homotypic synonym: \equiv *Alchemilla arvensis* subsp. *maroccana* (Hyl. & Rothm.) Maire in Emb. & Maire, Cat. Pl. Maroc 4: 1023 (1941).

The species was described by Hylander (1938). Although the holotype sheet has the handwritten label ‘*Alchemilla maroccana* Hyl. et Rothm. n. sp.’ it was finally described as *Aphanes*. Maire combined it into *Alchemilla* as a subspecies (Emberger & Maire, 1941).

IPNI, WFO and POWO (2024) record as an accepted name ‘*Alchemilla maroccana* (Hyl. & Rothm.) Devesa, Fl. Vasc. Andalucía Occid. 2: 37 (1987)’, however, this combination was not made in that work (cf. Devesa, 1987 and pers. comm.). Neither did Maire (cf. Emberger & Maire, 1941) make the combination ‘*Aphanes arvensis* subsp. *maroccana* (Hyl. & Rothm.) Maire in Jahand. & Maire, Cat. Pl. Maroc 4: 1023 (1941)’ mentioned by POWO and WFO.

The transfer as a species to the genus *Alchemilla* is proposed here, together with the correction of incorrectly attributed names.

This species occurs in Morocco and CW and SW Iberian Peninsula (Devesa, 1986; Frost-Olsen, 1998).

Conflict of interest

None.

References

- Christenhusz, M.J.M. & Väre, H. (2012). New combinations in *Potentilla* (Rosaceae) for the Nordic Flora. *Phytotaxa*, 57, 1-5. <https://doi.org/10.11646/phytotaxa.57.1.1>
- Devesa, J.A. (1986). 144. *Aphanes floribunda* (Murb.) Rothm. y *Aphanes maroccana* Hylander & Rothm. en Andalucía Occidental. *Lagascalia*, 14(1), 74-76.
- Devesa, J.A. (1987). ROSACEAE. In B. Valdés, S. Talavera & F. Galiano (Eds.), *Flora vascular de Andalucía Occidental*. Vol. 2 (pp. 23-43). Barcelona: Ketres.
- Dobeš, Ch. & Paule, J. (2010). A comprehensive chloroplast DNA-based phylogeny of the genus *Potentilla* (Rosaceae): implications for its geographic origin, phylogeography and generic circumscription. *Molecular Phylogenetics and Evolution*, 56, 156-175. <https://doi.org/10.1016/j.ympev.2010.03.005>
- Emberger, L. & Maire, R. (1941). *Catalogue des plantes du Maroc*. Volume IV. Supplément aux volumes I, II et III. Alger: Imprimerie Minerva.
- Eriksson, T., Persson, N.L. & Smedmark, J.E.E. (2022). What is *Potentilla*? A phylogeny-based taxonomy for Potentillinae (Rosaceae). *Taxon*, 71(3), 493-505. <https://doi.org/10.1002/tax.12679>
- Frost-Olsen, P. (1997). A new species of *Aphanes* L. (Rosaceae) from Portugal. *Anales del Jardín Botánico de Madrid*, 55(1), 195-197.
- Frost-Olsen, P. (1998). *Aphanes* L. In F. Muñoz Garmendia & C. Navarro (Eds.), *Flora Iberica: plantas vasculares de la Península Ibérica e Islas Baleares*. Vol. VI. Rosaceae (pp. 357-369). Madrid: Real Jardín Botánico. CSIC.
- Gehrke, B., Bräuchler, C., Romoleroux, K., Lundberg, M., Heubl, G. & Eriksson, T. (2008). Molecular phylogenetics of *Alchemilla*, *Aphanes* and *Lachemilla* (Rosaceae) inferred from plastid and nuclear intron and spacer DNA sequences, with comments on generic classification. *Molecular phylogenetics and evolution*, 47(3), 1030-1044. <https://doi.org/10.1016/j.ympev.2008.03.004>
- Hylander, N. (1938). Eine neue *Aphanes*-art aus Marokko. *Svensk botanisk tidskrift*, 32(2), 186-190.
- IPNI (2024). *International Plant Names Index*. Published on the Internet <http://www.ipni.org>, The Royal Botanic Gardens, Kew, Harvard University Herbaria & Libraries and Australian National Herbarium. Retrieved 29 September 2024.

- Jahandiez, É. & Maire, R. (1932). *Catalogue des plantes du Maroc*. Tome Deuxième. Dicotylédones Archichlamydées. Alger: Imprimerie Minerva.
- Kalkman, C. (2004). Rosaceae. In K. Kubitzki (Ed.), *The families and genera of vascular plants, vol. 6. Dicotyledons: Celastrales, Oxalidales, Rosales, Cornales, Ericales* (pp. 343-386). Berlin: Springer.
- Lippert, W. (1984). Zur Kenntnis des *Aphanes microcarpa* Komplexes. *Mitteilungen der Botanischen Staatssammlung München*, 20, 451-464.
- Maire, R. (1980). *Flore de l'Afrique du Nord*. Volume XV. DICOTYLEDONAE: Rosales: Saxifragaceae, Pittosporaceae, Platanaceae, Rosaceae. Paris: Paul Lechevalier.
- Murbeck, S. (1915). Zur Morphologie und Systematik der Gattung *Alchemilla*. *Acta Universitatis Lundensis. Nova series*, 11(8), 1-17.
- Potter, D., Eriksson, T., Evans, R.C., Oh, S., Smedmark, J.E.E., Morgan, D.R., Kerr, M., Robertson, K.R., Arsenault, M., Dickinson, T.A. & Campbell, C.S. (2007). Phylogeny and classification of Rosaceae. *Plant Systematics and Evolution*, 266(1), 5-43. <http://dx.doi.org/10.1007/s00606-007-0539-9>
- POWO (2024). *Plants of the World Online*. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; <https://powo.science.kew.org/>. Retrieved 06 October 2024.
- Rothmaler, W. (1935). Systematische Vorarbeiten zu einer Monographie der Gattung *Alchemilla* (L.) Scop. emend. II. Die systematische Gliederung der Gattung. III. Notizen über das Subgenus *Aphanes* (L.). *Repertorium specierum novarum regni vegetabilis*, 38, 33-43.
- Stevens, P.F. (2001 onwards). *Angiosperm Phylogeny Website*. Version 14, July 2017 [and more or less continuously updated since]. will do. <http://www.mobot.org/MOBOT/research/APweb/>. Accessed on 06 October 2024.
- Walters, S.M. (1968). 24. *Aphanes* L. In T.G. Tutin, V.H. Heywood, N.A. Burges, D.M. Moore, D.H. Valentine, S.M. Walters & D.A. Webb (Eds.), *Flora Europaea Volume 2: Rosaceae to Umbelliferae* (p. 64). Cambridge: Cambridge University Press.
- WFO (2024). *World Flora Online*. Published on the Internet; <http://www.worldfloraonline.org>. Accessed on 06 October 2024.